

Publications Günther Schütz – Impact Factor \approx 10 plus

Status January 2012

Blood

Baumann, S., Dostert, A., Novac, N., Bauer, A., Schmid, W., Fas, S. C., Krueger, A., Heinzl, T., Kirchhoff, S., Schütz, G., and Krammer, P. H. (2005). Glucocorticoids inhibit activation-induced cell death (AICD) via direct DNA-dependent repression of the CD95-ligand gene by a glucocorticoid receptor dimer. **Blood** 106, 617-625.

Cell

Axel, R., Feigelson, P., and Schutz, G. (1976). Analysis of the complexity and diversity of mRNA from chicken liver and oviduct. **Cell** 7, 247-254.

Hynes, N. E., Groner, B., Sippel, A. E., Nguyen-Huu, M. C., and Schutz, G. (1977). mRNA complexity and egg white protein mRNA content in mature and hormone-withdrawn oviduct. **Cell** 11, 923-932.

Grez, M., Land, H., Giesecke, K., Schutz, G., Jung, A., and Sippel, A. E. (1981). Multiple mRNAs are generated from the chicken lysozyme gene. **Cell** 25, 743-752.

Renkawitz, R., Beug, H., Graf, T., Matthias, P., Grez, M., and Schutz, G. (1982). Expression of a chicken lysozyme recombinant gene is regulated by progesterone and dexamethasone after microinjection into oviduct cells. **Cell** 31, 167-176.

Renkawitz, R., Schutz, G., von der Ahe, D., and Beato, M. (1984). Sequences in the promoter region of the chicken lysozyme gene required for steroid regulation and receptor binding. **Cell** 37, 503-510.

Miksicek, R., Heber, A., Schmid, W., Danesch, U., Posseckert, G., Beato, M., and Schutz, G. (1986). Glucocorticoid responsiveness of the transcriptional enhancer of Moloney murine sarcoma virus. **Cell** 46, 283-290.

Becker, P. B., Ruppert, S., and Schutz, G. (1987). Genomic footprinting reveals cell type-specific DNA binding of ubiquitous factors. **Cell** 51, 435-443.

Jantzen, H. M., Strahle, U., Gloss, B., Stewart, F., Schmid, W., Boshart, M., Miksicek, R., and Schutz, G. (1987). Cooperativity of glucocorticoid response elements located far upstream of the tyrosine aminotransferase gene. **Cell** 49, 29-38.

Stewart, A. F., and Schutz, G. (1987). Camptothecin-induced in vivo topoisomerase I cleavages in the transcriptionally active tyrosine aminotransferase gene. **Cell** 50, 1109-1117.

Boshart, M., Weih, F., Schmidt, A., Fournier, R. E., and Schutz, G. (1990). A cyclic AMP response element mediates repression of tyrosine aminotransferase gene transcription by the tissue-specific extinguisher locus Tse-1. **Cell** 61, 905-916.

Ruppert, S., Boshart, M., Bosch, F. X., Schmid, W., Fournier, R. E., and Schutz, G. (1990). Two genetically defined trans-acting loci coordinately regulate overlapping sets of liver-specific genes. **Cell** 61, 895-904.

Boshart, M., Weih, F., Nichols, M., and Schutz, G. (1991). The tissue-specific extinguisher locus TSE1 encodes a regulatory subunit of cAMP-dependent protein kinase. **Cell** 66, 849-859.

Bourtchuladze, R., Frenguelli, B., Blendy, J., Cioffi, D., Schutz, G., and Silva, A. J. (1994). Deficient long-term memory in mice with a targeted mutation of the cAMP-responsive element-binding protein. **Cell** 79, 59-68.

Beato, M., Herrlich, P., and Schütz, G. (1995). Steroid hormone receptors: many actors in search of a plot. **Cell** 83, 851-857.

Mangelsdorf, D. J., Thummel, C., Beato, M., Herrlich, P., Schutz, G., Umesono, K., Blumberg, B., Kastner, P., Mark, M., Chambon, P., and et al. (1995). The nuclear receptor superfamily: the second decade. **Cell** 83, 835-839.

Reichardt, H. M., Kaestner, K. H., Wessely, O., Tuckermann, J., Angel, P., Kretz, O., Bock, R., Schmid, W., Herrlich, P., and Schütz, G. (1998). DNA binding of the glucocorticoid receptor is not essential for survival. *Cell* **93**, 531-541.

Yadav, V. K., Ryu, J.-H., Suda, N., Tanaka, K., Gingrich, J. A., Schütz, G., Glorieux, F. H., Chiang, C. Y., Zajac, J. D., Insogna, K. L., Mann, J. J., Hen, R., Ducy, P., and Karsenty, G. (2008). Lrp5 controls bone formation by inhibiting serotonin synthesis in the duodenum. *Cell* **135**, 825-837.

Uhlenhaut, N. H., Jakob, S., Anlag, K., Eisenberger, T., Sekido, R., Kress, J., Treier, A.-C., Klugmann, C., Klasen, C., Holter, N. I., Riethmacher, D., Schütz, G., Cooney, A. J., Lovell-Badge, R., and Treier, M. (2009). Somatic sex reprogramming of adult ovaries to testes by FOXL2 ablation. *Cell* **139**, 1130-1142.

Cell Metabolism

Rauch, A., Seitz, S., Baschant, U., Schilling, A. F., Illing, A., Stride, B., Kirilov, M., Mandic, V., Takacz, A., Schmidt-Ullrich, R., Ostermay, S., Schinke, T., Spanbroek, R., Zaiss, M. M., Angel, P. E., Lerner, U. H., David, J.-P., Reichardt, H. M., Amling, M., Schütz, G., and Tuckermann, J. P. (2010). Glucocorticoids suppress bone formation by attenuating osteoblasts differentiation via the nonmreic glucocorticoid receptor. *Cell Metabolism* **11**, 517-531.

EMBO J.

Matthias, P. D., Renkawitz, R., Grez, M., and Schutz, G. (1982). Transient expression of the chicken lysozyme gene after transfer into human cells. *EMBO J* **1**, 1207-1212.

Schmid, W., Scherer, G., Danesch, U., Zentgraf, H., Matthias, P., Strange, C. M., Rowekamp, W., and Schutz, G. (1982). Isolation and characterization of the rat tryptophan oxygenase gene. *EMBO J* **1**, 1287-1293.

Matthias, P. D., Bernard, H. U., Scott, A., Brady, G., Hashimoto-Gotoh, T., and Schutz, G. (1983). A bovine papilloma virus vector with a dominant resistance marker replicates extrachromosomally in mouse and E. coli cells. *EMBO J* **2**, 1487-1492.

Becker, P., Renkawitz, R., and Schutz, G. (1984). Tissue-specific DNaseI hypersensitive sites in the 5'-flanking sequences of the tryptophan oxygenase and the tyrosine aminotransferase genes. *EMBO J* **3**, 2015-2020.

Brady, G., Funk, A., Mattern, J., Schutz, G., and Brown, R. (1985). Use of gene transfer and a novel cosmid rescue strategy to isolate transforming sequences. *EMBO J* **4**, 2583-2588.

Cato, A. C., Miksicek, R., Schutz, G., Arnemann, J., and Beato, M. (1986). The hormone regulatory element of mouse mammary tumour virus mediates progesterone induction. *EMBO J* **5**, 2237-2240.

Danesch, U., Gloss, B., Schmid, W., Schutz, G., Schule, R., and Renkawitz, R. (1987). Glucocorticoid induction of the rat tryptophan oxygenase gene is mediated by two widely separated glucocorticoid-responsive elements. *EMBO J* **6**, 625-630.

Muller, G., Ruppert, S., Schmid, E., and Schutz, G. (1988). Functional analysis of alternatively spliced tyrosinase gene transcripts. *EMBO J* **7**, 2723-2730.

Ruppert, S., Muller, G., Kwon, B., and Schutz, G. (1988). Multiple transcripts of the mouse tyrosinase gene are generated by alternative splicing. *EMBO J* **7**, 2715-2722.

Strahle, U., Schmid, W., and Schutz, G. (1988). Synergistic action of the glucocorticoid receptor with transcription factors. *EMBO J* **7**, 3389-3395.

Schmid, W., Strahle, U., Schutz, G., Schmitt, J., and Stunnenberg, H. (1989). Glucocorticoid receptor binds cooperatively to adjacent recognition sites. *EMBO J* **8**, 2257-2263.

Beermann, F., Ruppert, S., Hummler, E., Bosch, F. X., Müller, G., Rüther, U., and Schütz, G. (1990). Rescue of the albino phenotype by introduction of a functional tyrosinase gene into mice. *EMBO J* **9**, 137-150.

Reik, A., Schutz, G., and Stewart, A. F. (1991). Glucocorticoids are required for establishment and maintenance of an alteration in chromatin structure: induction leads to a reversible disruption of nucleosomes over an enhancer. *EMBO J* **10**, 2569-2576.

Weih, F., Nitsch, D., Reik, A., Schutz, G., and Becker, P. B. (1991). Analysis of CpG methylation and genomic footprinting at the tyrosine aminotransferase gene: DNA methylation alone is not sufficient to prevent protein binding in vivo. **EMBO J** *10*, 2559-2567.

Nichols, M., Weih, F., Schmid, W., DeVack, C., Kowenz Leutz, E., Luckow, B., Boshart, M., and Schutz, G. (1992). Phosphorylation of CREB affects its binding to high and low affinity sites: implications for cAMP induced gene transcription. **EMBO J** *11*, 3337-3346.

Ruppert, S., Cole, T. J., Boshart, M., Schmid, E., and Schutz, G. (1992a). Multiple mRNA isoforms of the transcription activator protein CREB: generation by alternative splicing and specific expression in primary spermatocytes. **EMBO J** *11*, 1503-1512.

Ganss, R., Montoliu, L., Monaghan, A. P., and Schutz, G. (1994). A cell-specific enhancer far upstream of the mouse tyrosinase gene confers high level and copy number-related expression in transgenic mice. **EMBO J** *13*, 3083-3093.

Blendy, J. A., Kaestner, K. H., Schmid, W., Gass, P., and Schütz, G. (1996a). Targeting of the CREB gene leads to up-regulation of a novel CREB mRNA isoform. **EMBO J** *15*, 1098-1106.

Montoliu, L., Umland, T., and Schutz, G. (1996). A locus control region at -12 kb of the tyrosinase gene. **EMBO J** *15*, 6026-6034.

Hiemisch, H., Schutz, G., and Kaestner, K. H. (1997). Transcriptional regulation in endoderm development: characterization of an enhancer controlling Hnf3g expression by transgenesis and targeted mutagenesis. **EMBO J** *16*, 3995-4006.

Le Minh, N., Damiola, F., Tronche, F., Schütz, G., and Schibler, U. (2001). Glucocorticoid signaling inhibits food-induced phase-shifting of peripheral circadian oscillators. **EMBO J** *20*, 7128-7136.

Reichardt, H. M., Tuckermann, J. P., Göttlicher, M., Vujic, M., Weih, F., Angel, P., Herrlich, P., and Schütz, G. (2001). Repression of inflammatory responses in the absence of DNA-binding by the glucocorticoid receptor. **EMBO J** *20*, 7168-7173.

Behrens, A., Sibilina, M., David, J.-P., Schütz, G., and Wagner, E. F. (2002). Impaired postnatal hepatocyte proliferation and liver regeneration in mice lacking c-jun in the liver. **EMBO J** *21*, 1782-1790.

Li, N., Zhang, Y., Naylor, M. J., Schatzmann, F., Maurer, F., Wintermantel, T., Schütz, G., Müller, U., Streuli, C. H., and Hynes, N. E. (2005). b1 integrins regulate mammary gland proliferation and maintain the integrity of mammary alveoli. **EMBO J** *24*, 1942-1953.

Sutherland, K. D., Vaillant, F., Alexander, W. S., Wintermantel, T. M., Forrest, N. C., Holroyd, S. L., McManus, E. J., Schutz, G., Watson, C. J., Chodosh, L. A., Lindeman, G. J., and Visvader, J. E. (2006). c-myc as a mediator of accelerated apoptosis and involution in mammary glands lacking Socs3. **EMBO J** *25*, 5805-5815.

Genes Dev.

Weih, F., Stewart, A. F., Boshart, M., Nitsch, D., and Schutz, G. (1990). In vivo monitoring of a cAMP-stimulated DNA-binding activity. **Genes Dev** *4*, 1437-1449.

Ruppert, S., Kelsey, G., Schedl, A., Schmid, E., Thies, E., and Schutz, G. (1992b). Deficiency of an enzyme of tyrosine metabolism underlies altered gene expression in newborn liver of lethal albino mice. **Genes Dev** *6*, 1430-1443.

Kelsey, G., Ruppert, S., Beermann, F., Grund, C., Tanguay, R. M., and Schutz, G. (1993). Rescue of mice homozygous for lethal albino deletions: implications for an animal model for the human liver disease tyrosinemia type 1. **Genes Dev** *7*, 2285-2297.

Nitsch, D., Boshart, M., and Schutz, G. (1993a). Extinction of tyrosine aminotransferase gene activity in somatic cell hybrids involves modification and loss of several essential transcriptional activators. **Genes Dev** *7*, 308-319.

Cole, T. J., Blendy, J. A., Monaghan, A. P., Kriegstein, K., Schmid, W., Aguzzi, A., Fantuzzi, G., Hummler, E., Unsicker, K., and Schutz, G. (1995). Targeted disruption of the glucocorticoid receptor gene blocks adrenergic chromaffin cell development and severely retards lung maturation. **Genes Dev** *9*, 1608-1621.

Kaestner, K. H., Silberg, D. G., Traber, P. G., and Schutz, G. (1997). The mesenchymal winged helix transcription factor Fkh6 is required for the control of gastrointestinal proliferation and differentiation. **Genes Dev** *11*, 1583-1595.

Bauer, A., Tronche, F., Wessely, O., Kellendonk, C., Reichardt, H. M., Steinlein, P., Schütz, G., and Beug, H. (1999). The glucocorticoid receptor is required for stress erythropoiesis. **Genes Dev** *13*, 2996-3002.

Kaestner, K. H., Katz, J., Liu, Y., Drucker, D., and Schütz, G. (1999). Inactivation of the winged helix transcription factor HNF3a affects glucose homeostasis and islet glucagon gene expression in vivo. **Genes Dev** *13*, 495-504.

Tronche, F., Opherk, C., Moriggl, R., Kellendonk, C., Reimann, A., Schwake, L., Reichardt, H. M., Stangl, K., Gau, D., Hoeflich, A., Beug, H., Schmid, W., and Schütz, G. (2004). Glucocorticoid receptor function in hepatocytes is essential to promote post-natal body growth. **Genes Dev** *18*, 492-497.

Engblom, D., Kornfeld, J.-W., Schwake, L., Tronche, F., Reimann, A., Beug, H., Hennighausen, L., Moriggl, R., and Schütz, G. (2007). Direct glucocorticoid receptor-Stat5 interaction in hepatocytes controls body size and maturation-related gene expression. **Genes Dev** *21*, 1157-1162.

Liu, H.-K., Belz, T., Bock, D., Takacs, A., Wu, H., Lichter, P., Chai, M., and Schütz, G. (2008). The nuclear receptor tailless is required for neurogenesis in the adult subventricular zone. **Genes Dev** *22*, 2473-2478.

Liu, H.-K., Wang, Y., Belz, T., Bock, D., Takacs, A., Radlwimmer, B., Barbus, S., Reifenberger, G., Lichter, P., and Schütz, G. (2010). The nuclear receptor tailless induces neural stem cell expansion and brain tumor initiation. **Genes Dev** *24*, 683-695.

Oury, F., Yadav, V.K., Wang, Y., Zhou, B., Liu, X.S., Guo, X.E., Tecott, L.H., Schütz, G. Means, A.B., and Karsenty, G. (2010). CREB mediates brain serotonin regulation of bone mass through its expression in ventromedial hypothalamic neurons. **Genes Dev** *24*, 2330-2342.

J. Cell Biol.

Tuckermann, J. P., Reichardt, H. M., Arribas, R., Richter, K. H., Schütz, G., and Angel, P. (1999). The DNA-binding-independent function of the glucocorticoid receptor mediates repression of AP-1-dependent genes in skin. **J Cell Biol** *147*, 1-6.

Naylor, M. J., Li, N., Cheung, J., Lowe, E. T., Lambert, E., Marlow, R., Wang, P., Schatzmann, F., Wintermantel, T., Schütz, G., Clarke, A. R., Müller, U., Hynes, N. E., and Streuli, C. H. (2005). Ablation of b1 integrin in mammary epithelium reveals a key role for integrin in glandular morphogenesis and differentiation. **J Cell Biol** *171*, 717-728.

J. Clin. Invest.

Limbourg, F. P., Huang, Z., Plumier, J.-C., Simoncini, T., Fujioka, M., Tuckermann, J., Schütz, G., Moskowitz, M. A., and Liao, J. K. (2002). Enhancement of cerebral blood flow and stroke protection mediated by non-nuclear actions of the glucocorticoid receptor. **J Clin Invest** *110*, 1729-1738.

Kero, J., Ahmed, K., Wettschureck, N., Tunaru, S., Wintermantel, T., Greiner, E., Schütz, G., and Offermanns, S. (2007). Thyrocyte-specific G_q/G₁₁ deficiency impairs thyroid function and goiter development. **J Clin Invest** *117*, 2399-2407.

Tuckermann, J. P., Kleiman, A., Moriggl, R., Spanbroek, R., Neumann, A., Illing, A., Clausen, B. E., Stride, B., Forster, I., Habenicht, A. J., Reichardt, H. M., Tronche, F., Schmid, W., and Schutz, G. (2007). Macrophages and neutrophils are the targets for immune suppression by glucocorticoids in contact allergy. **J Clin Invest** *117*, 1381-1390.

Grosse-Wilde, A., Voloshanenko, O., Bailey, S. L., Longton, G. M., Schaefer, U., Csernok, A. I., Schütz, G., Greiner, E. F., Kemp, C. J., and Walczak, H. (2008). TRAIL-R deficiency enhances lymph node metastasis without affecting primary tumor development. **J Clin Invest** *118*, 100-110.

Usher, M. G., Duan, S. Z., Ivashchenko, C., Frieler, R., Berger, S., Schütz, G., Lumeng, C., and Mortensen, R. M. (2010). Myeloid mineralocorticoid receptor controls macrophage polarization and cardiovascular hypertrophy and remodelling in mice. **J Clin Invest** *120*, 3350-3364.

Mol. Cell

Yuan, X., Zhou, Y., Casanova, E., Kenzelmann, M., Kiss, E., Gretz, N., Gröne, H.-J., Schütz, G., and Grummt, I. (2005). Nucleolar disruption by inactivation of the RNA polymerase I transcription factor TIF-IA leads to cell cycle arrest and p53-mediated apoptosis. **Mol Cell** *19*, 77-87.

Nat. Genet.

Tronche, F., Kellendonk, C., Kretz, O., Gass, P., Anlag, K., Orban, P. C., Bock, R., Klein, R., and Schütz, G. (1999). Disruption of the glucocorticoid receptor gene in the nervous system results in reduced anxiety. **Nat Genet** *23*, 99-103.

Mantamadiotis, T., Lemberger, T., Bleckmann, S., Kern, H., Kretz, O., Villalba-Martin, A., Tronche, F., Kellendonk, C., Gau, D., Kapfhammer, J., Otto, C., Schmid, W., and Schütz, G. (2002). Disruption of CREB function in brain leads to neurodegeneration. **Nat Genet** *31*, 47 - 54.

Nat. Med.

Wirth, A., Benyó, Z., Lukasowa, M., Leutgeb, B., Wettschureck, N., Gorbey, S., Horváth, B., Greiner, E., Lemmer, B., Schütz, G., Gutkind, S., and Offermanns, S. (2008). G₁₂/G₁₃-LARG-mediated signalling in vascular smooth muscle is required for salt-induced hypertension. **Nat Med** *14*, 64-68.

Nat. Neurosci.

Karst, H., Karten, Y. J. G., Reichardt, H. M., de Kloet, E. R., Schütz, G., and Joels, M. (2000). Corticosteroid actions in hippocampus require DNA-binding of glucocorticoid receptors. **Nat Neurosci** *3*, 977-978.

Ramanan, N., Shen, Y., Sarsfield, S. L., Lemberger, T., Schütz, G., Linden, D. J., and Ginty, D. D. (2005). SRF mediates activity-induced gene expression and synaptic plasticity but not neuronal viability. **Nat Neurosci** *8*, 759-767.

Knöll, B., Kretz, O., Fiedler, C., Alberti, S., Schütz, G., Frotscher, M., and Nordheim, A. (2006). Serum response factor controls neuronal circuit assembly in the hippocampus. **Nat Neurosci** *9*, 195-204.

Ambroggi, F., Turiault, M., Milet, A., Deroche-Gamonet, V., Parnaudeau, S., Balado, E., Barik, J., van der Veen, R., Maroteaux, G., Lemberger, T., Schütz, G., Lazar, M., Marinelli, M., Piazza, P. V., and Tronche, F. (2009). Stress and addiction: a specific neuronal cell type for GR-induced facilitation of cocaine seeking. **Nat Neurosci** *12*, 247-249.

Nature

Groner, B., Hynes, N., Sippel, A., and Schutz, G. (1976). Induction of specific proteins in hyphae of *Achlya ambisexualis* by the steroid hormone antheridiol. **Nature** *261*, 599-601.

Land, H., Schutz, G., Schmale, H., and Richter, D. (1982). Nucleotide sequence of cloned cDNA encoding bovine arginine vasopressin-neurophysin II precursor. **Nature** *295*, 299-303.

Land, H., Grez, M., Ruppert, S., Schmale, H., Rehbein, M., Richter, D., and Schutz, G. (1983). Deduced amino acid sequence from the bovine oxytocin-neurophysin I precursor cDNA. **Nature** *302*, 342-344.

Ruppert, S., Scherer, G., and Schutz, G. (1984). Recent gene conversion involving bovine vasopressin and oxytocin precursor genes suggested by nucleotide sequence. **Nature** 308, 554-557.

von der Ahe, D., Janich, S., Scheidereit, C., Renkawitz, R., Schutz, G., and Beato, M. (1985). Glucocorticoid and progesterone receptors bind to the same sites in two hormonally regulated promoters. **Nature** 313, 706-709.

Becker, P. B., Gloss, B., Schmid, W., Strahle, U., and Schutz, G. (1986). In vivo protein-DNA interactions in a glucocorticoid response element require the presence of the hormone. **Nature** 324, 686-688.

Klock, G., Strahle, U., and Schutz, G. (1987). Oestrogen and glucocorticoid responsive elements are closely related but distinct. **Nature** 329, 734-736.

Strahle, U., Boshart, M., Klock, G., Stewart, F., and Schutz, G. (1989). Glucocorticoid- and progesterone-specific effects are determined by differential expression of the respective hormone receptors. **Nature** 339, 629-632.

Schedl, A., Montoliu, L., Kelsey, G., and Schutz, G. (1993). A yeast artificial chromosome covering the tyrosinase gene confers copy number-dependent expression in transgenic mice. **Nature** 362, 258-261.

Blendy, J. A., Kaestner, K. H., Weinbauer, G. F., Nieschlag, E., and Schütz, G. (1996b). Severe impairment of spermatogenesis in mice lacking the CREM gene. **Nature** 380, 162-165.

Monaghan, A. P., Bock, D., Gass, P., Schwäger, A., Wolfner, D. P., Lipp, H.-P., and Schütz, G. (1997). Defective limbic system in mice lacking the tailless gene. **Nature** 390, 515-517.

Herzig, S., Fanxin, L., Jhala, U., Hedrick, S., Quinn, R., Bauer, A., Schütz, G., Yoon, C., Puisgever, P., Spiegelman, B., and Montminy, M. (2001). CREB regulates hepatic gluconeogenesis through the coactivator PGC-1. **Nature** 413, 179-183.

Neuron

Gau, D., Lemberger, T., von Gall, C., Kretz, O., Le Minh, N., Gass, P., Schmid, W., Schibler, U., Korf, H. W., and Schütz, G. (2002). Phosphorylation of CREB Ser-142 regulates light-induced phase shifts of the circadian clock. **Neuron** 34, 245-253.

Wintermantel, T., Campbell, R. E., Porteous, R., Bock, D., Gröne, H.-J., Todman, M. G., Korach, K. S., Greiner, E., Pérez, C. A., Schütz, G., and Herbison, A. E. (2006). Definition of estrogen receptor pathway critical for estrogen positive feedback to gonadotropin-releasing hormone neurons and fertility. **Neuron** 52, 271-280.

Zhang, S. J., Steijaert, M. N., Lau, D., Schutz, G., Delucinge-Vivier, C., Descombes, P., and Bading, H. (2007). Decoding NMDA Receptor Signaling: Identification of Genomic Programs Specifying Neuronal Survival and Death. **Neuron** 53, 549-562.

Engblom, D., Bilbao-Leis, A., Sanchis-Segura, C., Dahan, L., Perreau-Lenz, S., Baland, B., Rodriguez Parkitna, J., Lujan, R., Halbout, B., Mameli, M., Parlato, R., Sprengel, R., Lüscher, C., Schütz, G., and Spanagel, R. (2008). Glutamate receptors on dopamine neurons control the persistence of cocaine-seeking. **Neuron** 59, 497-508.

PNAS

Ruiz-Carrilo, A., Beato, M., Schutz, G., Feigelson, P., and Allfrey, V. G. (1973). Cell-free translation of the globin message within polydisperse high-molecular-weight ribonucleic acid of avian erythrocytes. **Proc Natl Acad Sci U S A** 70, 3641-3645.

Schutz, G., Beato, M., and Feigelson, P. (1973). Messenger RNA for hepatic tryptophan oxygenase: its partial purification, its translation in a heterologous cell-free system, and its control by glucocorticoid hormones. **Proc Natl Acad Sci U S A** 70, 1218-1221.

Sippel, A. E., Stavrianopoulos, J. G., Schutz, G., and Feigelson, P. (1974). Translational properties of rabbit globin mRNA after specific removal of poly(A) with ribonuclease H. **Proc Natl Acad Sci U S A** 71, 4635-4639.

Killewich, L., Schutz, G., and Feigelson, P. (1975). Functional level of rat liver tryptophan 2,3-dioxygenase messenger RNA during superinduction of enzyme with actinomycin D. **Proc Natl Acad Sci U S A** 72, 4285-4287.

Schutz, G., Killewich, L., Chen, G., and Feigelson, P. (1975). Control of the mRNA for hepatic tryptophan oxygenase during hormonal and substrate induction. **Proc Natl Acad Sci U S A** 72, 1017-1020.

Nguyen-Huu, M. C., Sippel, A. A., Hynes, N. E., Groner, B., and Schutz, G. (1978). Preferential transcription of the ovalbumin gene in isolated hen oviduct nuclei by RNA polymerase B. **Proc Natl Acad Sci U S A** 75, 686-690.

Lindenmaier, W., Nguyen-Huu, M. C., Lurz, R., Stratmann, M., Blin, N., Wurtz, T., Hauser, H. J., Sippel, A. E., and Schutz, G. (1979). Arrangement of coding and intervening sequences of chicken lysozyme gene. **Proc Natl Acad Sci U S A** 76, 6196-6200.

Nguyen-Huu, M. C., Stratmann, M., Groner, B., Wurtz, T., Land, H., Giesecke, K., Sippel, A. E., and Schutz, G. (1979). Chicken lysozyme gene contains several intervening sequences. **Proc Natl Acad Sci U S A** 76, 76-80.

Jung, A., Sippel, A. E., Grez, M., and Schutz, G. (1980). Exons encode functional and structural units of chicken lysozyme. **Proc Natl Acad Sci U S A** 77, 5759-5763.

Scherer, G., Schmid, W., Strange, C. M., Rowekamp, W., and Schutz, G. (1982). Isolation of cDNA clones coding for rat tyrosine aminotransferase. **Proc Natl Acad Sci U S A** 79, 7205-7208.

Hashimoto, S., Schmid, W., and Schutz, G. (1984). Transcriptional activation of the rat liver tyrosine aminotransferase gene by cAMP. **Proc Natl Acad Sci U S A** 81, 6637-6641.

Shinomiya, T., Scherer, G., Schmid, W., Zentgraf, H., and Schutz, G. (1984). Isolation and characterization of the rat tyrosine aminotransferase gene. **Proc Natl Acad Sci U S A** 81, 1346-1350.

Schmid, W., Muller, G., Schutz, G., and Gluecksohn-Waelsch, S. (1985). Deletions near the albino locus on chromosome 7 of the mouse affect the level of tyrosine aminotransferase mRNA. **Proc Natl Acad Sci U S A** 82, 2866-2869.

Strahle, U., Klock, G., and Schutz, G. (1987). A DNA sequence of 15 base pairs is sufficient to mediate both glucocorticoid and progesterone induction of gene expression. **Proc Natl Acad Sci U S A** 84, 7871-7875.

Ankenbauer, W., Strahle, U., and Schutz, G. (1988). Synergistic action of glucocorticoid and estradiol responsive elements. **Proc Natl Acad Sci U S A** 85, 7526-7530.

Kluppel, M., Beermann, F., Ruppert, S., Schmid, E., Hummler, E., and Schutz, G. (1991). The mouse tyrosinase promoter is sufficient for expression in melanocytes and in the pigmented epithelium of the retina. **Proc Natl Acad Sci U S A** 88, 3777-3781.

Beermann, F., Schmid, E., and Schütz, G. (1992). Expression of the mouse tyrosinase gene during embryonic development: recapitulation of the temporal regulation in transgenic mice. **Proc Natl Acad Sci U S A** 89, 2809-2813.

Strahle, U., Schmidt, A., Kelsey, G., Stewart, A. F., Cole, T. J., Schmid, W., and Schutz, G. (1992). At least three promoters direct expression of the mouse glucocorticoid receptor gene. **Proc Natl Acad Sci U S A** 89, 6731-6735.

Kaestner, K. H., Lee, K. H., Schlondorff, J., Hiemisch, H., Monaghan, A. P., and Schutz, G. (1993). Six members of the mouse forkhead gene family are developmentally regulated. **Proc Natl Acad Sci U S A** 90, 7628-7631.

Nitsch, D., Boshart, M., and Schutz, G. (1993b). Activation of the tyrosine aminotransferase gene is dependent on synergy between liver-specific and hormone-responsive elements. **Proc Natl Acad Sci U S A** 90, 5479-5483.

Hummler, E., Cole, T. J., Blendy, J. A., Ganss, R., Aguzzi, A., Schmid, W., Beermann, F., and Schutz, G. (1994). Targeted mutation of the CREB gene: compensation within the CREB/ATF family of transcription factors. **Proc Natl Acad Sci U S A** 91, 5647-5651.

Montoliu, L., Blendy, J. A., Cole, T. J., and Schutz, G. (1995). Analysis of perinatal gene expression: hormone response elements mediate activation of a lacZ reporter gene in liver of transgenic mice. **Proc Natl Acad Sci U S A** 92, 4244-4248.

Berger, S., M., B., Schmid, W., Cole, T. J., Peters, J., Watanabe, H., Kriz, W., Warth, R., Greger, R., and Schütz, G. (1998). Mineralocorticoid receptor knock out mice: Pathophysiology of Na⁺ metabolism. **Proc Natl Acad Sci U S A** 95, 9424-9429.

Rudolph, D., Tafuri, A., Gass, P., Hämmerling, G. J., Arnold, B., and Schütz, G. (1998). Impaired fetal T-cell development and perinatal lethality in mice lacking the cAMP response element binding protein (CREB). **Proc Natl Acad Sci U S A** *95*, 4481-4486.

Alberti, S., Krause, S. M., Kretz, O., Philippar, U., Lemberger, T., Casanova, E., Wiebel, F., Schwarz, H., Frotscher, M., Schütz, G., and Nordheim, A. (2005). Neuronal migration in the murine rostral migratory stream requires SRF. **Proc Natl Acad Sci U S A** *102*, 6148-6153.

Karst, H., Berger, S., Turiault, M., Tronche, F., Schütz, G., and Joels, M. (2005). Mineralocorticoid receptors are indispensable for nongenomic modulation of hippocampal glutamate transmission by corticosterone. **Proc Natl Acad Sci U S A** *102*, 19204-19207.

Berger, S., Wolfer, D. P., Selbach, O., Alter, H., Erdmann, G., Reichardt, H. M., Chepkova, A. N., Welzl, H., Haas, H. L., Lipp, H.-P., and Schütz, G. (2006). Loss of the limbic mineralocorticoid receptor impairs behavioral plasticity. **Proc Natl Acad Sci U S A** *103*, 195-200.

Gantois, I., Fang, K., Jiang, L., Babovic, D., Lawrence, A. J., Ferreri, V., Teper, Y., Jupp, B., Ziebell, J., Morganti-Kossmann, C. M., O'Brien, T. J., Nally, R., Schutz, G., Waddington, J., Egan, G. F., and Drago, J. (2007). Ablation of D1 dopamine receptor-expressing cells generates mice with seizures, dystonia, hyperactivity, and impaired oral behavior. **Proc Natl Acad Sci U S A** *104*, 4182-4187.

Kenzelmann, M., Maertens, S., Hergenahn, M., Kueffer, S., Hotz-Wagenblatt, A., Li, L., Wang, S., Ittrich, C., Lemberger, T., Arribas-Prat, R., Jonnakuty, S., Hollstein, M., Schmid, W., Gretz, N., Gröne, H.-J., and Schütz, G. (2007). Microarray analysis of newly synthesized RNA in cells and animals. **Proc Natl Acad Sci U S A** *104*, 6164-6169.

Bilbao, A., Rodriguez Parkitna, J., Engblom, D., Perreau-Lenz, S., Sanchis-Segura, C., Schneider, M., Konopka, W., Westphal, M., Breen, G., Desrivieres, S., Klugmann, M., Guindalini, C., Vallada, H., Laranjeira, R., Rodriguez de Fonseca, F., Schumann, G., Schütz, G., and Spanagel, R. (2008). Loss of the Ca²⁺/calmodulin-dependent protein kinase type IV in dopaminergic neurons enhances behavioral effects of cocaine. **Proc Natl Acad Sci USA** *105*, 17549-17554.

Karst, H., Berger, S., Erdmann, G., Schütz, G., and Joels, M. (2010). Metaplasticity in amygdalar responses to the stress hormone corticosterone. **Proc Natl Acad Sci USA** *107*, 14449-14454.

Fusco, S., Ripoli, C., Podda, M.V., Ranieri, S.C., Leone, L., Toietta, G., McBurney, M.W., Schütz, G., Riccio, A., Grassi, C., Galeotti, T., and Pani, G. (2012). A role for neuronal cAMP responsive-element binding (CREB)-1 in brain responses to calorie restriction. **Proc Natl Acad Sci USA** *109*, 621-626.

Science

Maldonado, R., Blendy, J. A., Tzavara, E., Gass, P., Roques, B. P., Hanoune, J., and Schutz, G. (1996). Reduction of morphine abstinence in mice with a mutation in the gene encoding CREB. **Science** *273*, 657-659.

Balsalobre, A., Brown, S. A., Marcacci, L., Tronche, F., Kellendonk, C., Reichardt, H. M., Schütz, G., and Schibler, U. (2000). Glucocorticoid hormones can reset circadian time in peripheral tissues but not in the suprachiasmatic nucleus. **Science** *289*, 2344-2347.

Marsicano, G., Goodenough, S., Monory, K., Hermann, H., Eder, M., Cannich, A., Azad, S. C., Cascio, M. G., Ortega Gutiérrez, S., van der Stelt, M., López-Rodríguez, M. L., Casanova, E., Schütz, G., Zieglgänsberger, W., Di Marzo, V., Behl, C., and Lutz, B. (2003). CB1 cannabinoid receptors and on-demand defense against excitotoxicity. **Science** *302*, 84-88.

Refojo, D., Schweizer, M., Kuehne, C., Ehrenberg, S., Thoeniger, C., Vogl, A.M., Dedic, N., Schumacher, M., von Wolff, G., Avrabos, C., Touma, C., Engblom, D., Schütz, G., Nave, K.-A., Eder, M., Wotjak, C.T., Sillaber, I., Holsboer, F., Wurst, W., and Deussing, J.M. (2011). Glutamatergic and dopaminergic neurons mediate anxiogenic and anxiolytic effects of CRHR1. **Science** *333*, 1903-1907. *Scienceexpress*/1 September 2011/ 10.1126/science.1202107.